

Determination of the level of pesticide residues in irrigation soil from Akko, Gombe State, Nigeria

Mohammad Bashir Sulaiman^{1*}, Aliyu Fadawa Tijjani², Abdullahi Muhammad Gimba³

¹Department of Chemistry, Northwest University Sokoto, Sokoto State, Nigeria

²Department of Science Laboratory Technology, Federal Polytechnic of Oil and Gas, Bonny Island, Nigeria

³Department of Science Laboratory Technology, Gombe State Polytechnic Bajoga, Gombe State, Nigeria

Corresponding Author's email: sulaiman.mbashir@nwus.edu.ng; sulaimanmbashir35@gmail.com

Available at: <http://acsigeria.org/publications>

ABSTRACT

This study determined pesticide residues levels in irrigation soil from Akko, Gombe State, Nigeria. A total of 54 soil samples were collected from three different irrigation farmlands, and the pesticide residues were extracted and analyzed using Agilent GC 7890B, coupled with MSD 5977A. The mean concentration of dichlorvos in the soil samples ranged from 0.00 - 0.089 mg·kg⁻¹, while the concentrations of dimethoate, lindane, aldrin, cypermethrin, and pyrazophos were below the detectable limit in all the studied samples. Paraquat dichloride ranged from 0.00 - 0.084 mg·kg⁻¹, while chlorpyrifos ranged from 0.00 - 0.079 mg·kg⁻¹, and p,p'-DDT ranged from 0.078 - 0.098 mg·kg⁻¹, while heptachlor concentration ranged from 0.00 - 0.086 mg·kg⁻¹ in the soil samples. The levels of cyhalothrin ranged from 0.00 - 0.095 mg·kg⁻¹, dieldrin ranged from 0.00 - 0.085 mg·kg⁻¹, while that of endrin ranged from 0.00 - 0.097 mg·kg⁻¹. The concentrations of endosulfan ranged from 0.00 - 0.106 mg·kg⁻¹. This considerable level of these pesticide residues may accumulate in food crops through the uptake of soil and have a negative impact on human health. However, monitoring and continuous stringent regulations should be imposed with regard to the usage of pesticides in soil and foodstuff for public health protection.

KEYWORDS: Determination, irrigation, pesticide residues, soil.

1. INTRODUCTION

Pesticides are substances commonly used in modern agricultural practices to protect crops from various pests and diseases.¹⁻³ Many farmers use pesticides in order to increase yield and to protect their farms from diseases. However, after pesticide application, residues often remain in the soil and crops even after harvest, eventually entering the food chain.^{4,5} Pesticides are toxic to humans both acute and chronic health effects depending on the quantity and types of exposure.⁶ Conversely, the use of pesticides has significantly contributed to the increase in global food production over the past several decades.⁷

Globally, approximately 3 million tons of pesticides are used each year, representing a market value of US\$40 billion.⁸ In the European Union, nearly 500 substances are approved for pesticide use, with annual sales averaging 374,000 tons between 2011 and 2016.⁹ Although pesticides enhance agricultural productivity and play a crucial economic role, their excessive use has raised significant environmental and public health concerns.¹⁰ The widespread issue of diffuse agrochemical pollution has emerged as a significant threat to soil health¹¹, potentially undermining multiple United Nations Sustainable Development Goals related to the soil environment.^{12,13}

Soil contamination poses significant threats to soil functions, biodiversity, and food safety, while also facilitating the off-site spread of pollutants through wind and water erosion. This dispersal can disrupt sink ecosystems and create additional exposure pathways for humans and non-target organisms to harmful contaminants.^{13,14} Despite the significant impacts of soil contamination, EU regulations do not require monitoring pesticide residues in soil, in contrast to the mandatory water monitoring established by the Water Framework Directive.⁷ There is a lack of extensive international studies on pesticide residues in soil, as existing research typically analyzes only individual pesticides or a small group of them.^{7,15,16}

Several studies have characterized the occurrence of both currently used and banned pesticides in soils at national or regional levels.¹⁷⁻²² Effective control of pesticide levels in food can only be achieved through systematic residue monitoring. In recent years, increased surveillance efforts have been implemented, emphasizing proper pesticide application rates and regulatory adherence.²³⁻²⁵ Despite the public health risks, the usage of these pesticides is increasing, and they are used for both agricultural and non-agricultural purposes in Nigeria.⁵ Hence, this study to investigate the level of pesticide residues in irrigation soil from Akko, Gombe State, Nigeria.

2. MATERIALS AND METHODS

2.1 Study area

The Akko Local Government Area is one of the eleven local governments in Gombe State. Akko is located between the latitudes of 10.97320° E and the longitudes of 10.28899° N of the equator. The area falls within the northern Guinea savanna zone of the Akko, Gombe State. Akko covers a total area of 2,627 km² (1,014 sq mi), with approximate population of 337,853.²⁶ The area receives an average annual rainfall of approximately 600 mm, which is sufficient for a single farming season. The annual rainfall pattern is erratic at the beginning of the rainy season, starting in April and intensifying as the season advances, raising the average from 600 to 1000 mm. The temperature as high as 41°C and the as low as 16°C. The generally high temperatures and low humidity favor high rates of evapotranspiration, leading to net water deficiencies. Akko L.G.A. is in the interior Savanna woodland, which can sustain large-scale livestock farming, as well as the cultivation of agricultural products like onions, spinach, tomatoes, rice, groundnuts, yams, cassava, and maize. The study area was selected due to dry farming activities.

2.2 Soil sample collection

Soil samples were collected from three farms (labelled Farm A, Farm B, and Farm C) using a nested sub-sampling approach. Each farm contains six sub-farms; from each sub-farm three soil cores were taken at a depth of 0–20 cm, yielding 18 samples per farm (3 samples × 6 sub-farms = 18). All 18 samples from each farm were composited into a single representative sample for that farm (i.e., one composite for Farm A, one for Farm B, and one for Farm C). Samples were labelled sequentially as A, B, and C corresponding to the three farms. The samples were all label and transported to laboratory for analysis.

2.3 Reagent

Reagents/Solvents: n-hexane and dichloromethane (HPLC grade). Copper powder and concentrated sulfuric acid were obtained from Loba Chemie India.

2.4 Sample Extraction

Five grams of the sample was weighed into a 50 mL centrifuge tube, and 30 mL aliquots of the extraction solvent consisting of a mixture of n-hexane/dichloromethane (1/1, v/v) were added and shaken for 15 min (using a universal shaker, IKA Werke). After 15 minutes, the sample mixture was ultrasonically extracted at 25°C for 20 min. The mixture was centrifuged at 8,000 rpm for 8 minutes, and the resulting supernatant was transferred into a separate container. The extraction process was repeated twice, after which all extracts were combined and concentrated using a rotary evaporator under mild vacuum until dryness. The residue was dissolved in 40 mL of n-hexane and analyzed using GC system: Agilent 7890A GC (Agilent Technologies)

2.5 Sample Clean up

Five grams of copper was added to the dissolved sample mixture, shaken for 5 min, and centrifuged at 8000 rpm for 5 min. The supernatant was collected, 3 mL of concentrated H₂SO₄ was added, and the mixture was vortex for 1 min. The mixture was again centrifuged at 2,000 rpm for 5 minutes, and the resulting supernatant was collected and treated with 1 mL of concentrated sulfuric acid. Vortex-mixed again for 1 min, and then centrifuged at 2000 r/min for 5 min. The supernatant solvent was dried by a rotary evaporator, and the residue was reconstituted with 1 mL n-hexane and then vortex-mixed.

2.6 Instrumental Analysis

A 1 µL aliquot of the reconstituted sample was injected into a gas chromatograph (Agilent 7890B GC coupled with a 5977A MSD) for analysis, using helium as the carrier gas. The column temperature program was set at 150 °C for 1 min, ramp at 25°C /min to 230 °C, hold for 2 min; and ramp at 20°C /min to 240 °C, hold for 1 min; ramp at 20 °C /min to 260 °C, and hold for 5 min; ramp at 3°C /min to 280 °C, and hold for 5 min; ramp at 2°C /min to 285 °C, and hold for 10 min. The injection port temperature and transfer line were set at 240 °C and 250 °C, respectively. The injection was made in the splitless mode with purge on after 0.75 min.²⁷

2.7 Preparation of Calibration Standards

Working standard solutions were prepared by diluting the stock solutions to 10 µg/mL in ethyl acetate. Appropriate aliquots were taken and further diluted with ethyl acetate to give a series of calibration standard solutions with concentrations of 10, 20, 50, 75, and 100 ng/mL. Extracted calibration

standards were prepared by spiking blank water samples (1 L) with the working standard solution prior to extraction to give concentrations of 10 ng/L, 20 ng/L, 10, 20, 50, 75, and 100 ng/L. After extraction and reconstitution (1 L to 1 mL), concentrations of these equate to of 0, 20, 50, 75, and 100 ng/L.²⁸

2.8 Method of Data Analysis

The residues were analyzed using a Shimadzu gas chromatograph (GC-2010), equipped with ⁶³Ni, an electron capture detector that allows the detection of contaminants even at trace concentrations. The GC conditions and detector response were adjusted to match the relative retention time and response. The capillary column was coated with ZB5 (30 m x 0.25 mm, 0.25 µm film thickness). Sample preparation, extraction, cleanup, and analysis were performed according to the procedure.²⁹

3. RESULTS AND DISCUSSION

3.1 Pesticide residues Percentage in the soil samples

The percentage of pesticide residues is presented in Figure 1. The percentage level of dichlorvos in the soil samples was in the range of 27.50% and 34.85% in farms B and C, respectively, while the percentage level of paraquat dichloride in the soil was 12.90% in farm A and below the detectable limit in farms B and C. The levels of chlorpyrifos in the soils were in the order of 24.38% and 32.78% in farms B and C, whereas they were below detectable in farm A. The p,p'-DDT distribution in the studied soils was 15.05%, 24.07%, and 32.36% in farms A, B, and C, respectively. The heptachlor in the soil was in the order of 13.21% in farm A, below detectable in farms B and C, while that of cyhalothrin had the following percentages in the same order: 14.59%, and dieldrin sulphate distribution in the studied soils was 13.05% and 24.07% in farms A and B, respectively. The percentage level of endrin in the soils was 14.90% in farm A and below the detectable limit in farms B and C, whereas that of endosulfan was 16.28% in similar orders.




Figure 1: Percentage (%) of pesticide residues in the soil samples

3.2 Levels of pesticide residues in soil samples

The pesticide levels in the soil samples are presented in Table 1. The mean concentration of dichlorvos in the soil samples ranged from 0.00 to 0.089 mg·kg⁻¹, and the concentrations of dimethoate, lindane, aldrin, cypermethrin, and pyrazophos in the samples were below the detectable limit. Paraquat dichloride concentration was 0.084 mg kg⁻¹ in farm A and was below the detectable limit in farms B and C. Chlorpyrifos concentration in the samples ranged from 0.00 to 0.079 mg·kg⁻¹, and concentrations of p,p'-DDT in the samples ranged from 0.078 to 0.098 mg·kg⁻¹, while heptachlor concentration ranged from 0.00 to 0.086 mg·kg⁻¹ in the samples. The levels of cyhalothrin in the samples ranged from 0.00 to 0.095 mg·kg⁻¹, the dieldrin ranged from 0.00 to 0.085 mg·kg⁻¹ in the sampled soils, while that of endrin ranged from 0.00 to 0.097 mg·kg⁻¹. The endosulfan concentrations in the samples ranged from 0.00 to 0.106 mg·kg⁻¹. Han et al. also reported that the content of chlorpyrifos, 7.2 - 77.2 µg/kg, was below the value of 0.079 mg kg⁻¹ obtained in this study³¹. The level

of p,p'-DDT obtained in this study was above (4.01 µg/kg) that reported in soils from the farms of Kumasi, Ghana.³⁰ and 34.0 reported from the soils of China.³¹

The level of cyhalothrin was higher than that reported by Han et al.³¹ but lower than that the cypermethrin reported by Han et al.³¹. Vincent et al. reported the detection of aldrin in soil 1.20 µg/kg, which is higher than the values obtained in this study.³⁰ In a Thailand project of food webs in rice paddies, a high level of aldrin was also detected in rice soils (28.6 µg/kg),³² while dieldrin values were reported as 1.5-3.9 µg/kg by Han et al., which were below the values of 0.078-0.085 mg·kg⁻¹ in this study.³¹ Moreover, this study showed that the levels of pesticide residues in the studied soils were higher than those in early reported studies conducted in other parts of the world.²²⁻³⁴ This could be due to unawareness and misuse of the use of OCP pesticides in Nigeria.⁷

Table 1: Levels of pesticide residues in soil (mg kg⁻¹) samples

S/No		Farm A	Farm B	Farm C
1	Dichlorvos	BDL	0.089±0.105	0.084±0.098
2	Dimethoate	BDL	BDL	BDL
3	Paraquat dichloride	0.084±0.105	BDL	BDL
4	Lindane	BDL	BDL	BDL
5	Chlorpyrifos	BDL	0.079±0.078	0.079±0.088
6	p,p'-DDT	0.098±0.089	0.078±0.098	0.078±0.100
7	Aldrin	BDL	BDL	BDL
8	Heptachlor	0.086±0.099	BDL	BDL
9	Cyhalothrin	0.095±0.110	BDL	BDL
10	Cypermethrin	BDL	BDL	BDL
11	Pyrazophos	BDL	BDL	BDL
12	Dieldrin	0.085±0.110	0.078±0.101	BDL
13	Endrin	0.097±0.974	BDL	BDL
14	Endosulfan	0.106±0.0970	BDL	BDL

BDL= below detectable limit

4. CONCLUSION

The present study was conducted to determine pesticide residues levels in the irrigation soil of Kunnuwal, Akko Local Government Area Gombe State, Nigeria. Generally, the pesticide residue concentrations in the various soils have been recorded at a reasonable concentration. Hence, the need for regular monitoring of pesticide residues should be encouraged, since an increase in misuse of pesticides for agricultural produce is still ongoing in Nigeria. Based on the findings of this study, the following recommendations for further action and studies are suggested: the present work critically focused on the pesticide residue assessment of irrigation soil in the study area; the results, therefore, suggest the need for further studies on agricultural produce to help in understanding the chemical form and behavior of the pesticide residues; there is a need to develop an environmental monitoring and management program for pesticide residues in Nigeria.

CONFLICT OF INTERESTS

The authors declare no conflict of interests.

REFERENCES

- (1) Guler, G.O.; Cakmak Y.S.; Dagli Z.; Aktumsek, A.; Ozparlak, H. Organo-chlorine pesticide residues in wheat from Konya region, Turkey. *Food Chem Toxicol.* **2010**, *4*, 1218-1221.
- (2) Ahoudi, H.; Gnandi, K.; Tanouayi, G.; Ouro-sama, K.; Yorke, J.C.; Creppy, E.E. and Moesch, C. Assessment of pesticides residues contents in the vegetables cultivated in urban area of

- Lome (southern Togo) and their risks on public health and the environment, Togo. *Int J Biol Chem Sci.* **2018**, *12*, 2172-2185.
- (3) Maigari, A.U.; Sulaiman, M.B.; Buhari, M.; Abdullahi, A.O. Pesticide residues in selected vegetables from Gombe markets, Gombe State, Nigeria: assessing the health impact. *Ife J Sci.* **2021**, *23*(1), 77-87.
- (4) Ononamadu, C.J.; Barau, M.M.; Salawu, K.; Ihegboro, G.O.; Owolafe, T.A.; Lawal, A.T.; Oshobu, M.L.; Unah, P.E. Screening of selected vegetables from Wudil, Farmlands in Kano State, Nigeria for organophosphorus and organochlorine pesticide residues. *J Environ Occup Sci.* **2019**, *82*, 20-25.
- (5) Sulaiman, M.B.; Maigari A.U.; Ihedioha, J.N.; Lawal, S.R.; Gimba A.M.; Shuhaibu, A.B. Levels and health risk assessment of organochlorine pesticide residues in vegetables from Yamaltu area in Gombe, Nigeria. *French-Ukraine Journal of Chemistry.* **2021**, *9*(1), 19-30.
- (6) Ahmad, M.F.; Ahmad, F.A.; Alsayegh, A.A.; Zeyaulah, M.; AlShahrani, A.M.; Muzammil K.; Saati, A.A.; Wahab, S.; Elbendary, E.Y.; Kambal, N.; Abdelrahman, M.H.; Hussain, S. Pesticides impacts on human health and the environment with their mechanisms of action and possible countermeasures. *Heliyon*, **2024**, *10*(7), e29128.
- (7) Silva, V.; Mol, H.G.J.; Zomer, P.; Tienstra, M.; Ritsema, C.J.; Geissen, V. Pesticide residues in European agricultural soils – A hidden reality unfolded. *Science of The Total Environment.* **2019**, *653*, 1532-1545. <https://doi.org/10.1016/j.scitotenv.2018.10.441>.
- (8) Pimentel, D. Pesticides and Pest Control. In: Peshin, R.; Dhawan, A.K. (eds) Integrated Pest Management: Innovation-Development Process. *Springer, Dordrecht.* **2009**. Doi; 10.1007/978-1-4020-8992-3_3
- (9) EC (2018) EU pesticides database - active substances. <http://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=activesubstance.selection&language=EN>. (Last access May) 2018.
- (10) Zhou, W.; Li, M.; Achal, V. A comprehensive review on environmental and human health impacts of chemical pesticide usage. *Emerging Contaminants.* **2025**, *11*, 1,100410.
- (11) Stolte, J.; Tesfai, M.; Oygarden, L.; Kvaerno, S.; Keizer, J.; Verheijen, F.; Panagos, P.; Ballabio, C.; Hessel, R. Soil threats in Europe: status, methods, drivers and effects on ecosystem services: deliverable 2.1 RECARE project. (98673 ed.) (JRC Technical reports). *European Commission DG Joint Research Centre.* **2016**. doi.org/10.2788/488054.
- (12) Keesstra, S. D.; Bouma, J.; Wallinga, J.; Tiftonell, P.; Smith, P.; Cerdà, A.; Montanarella, L.; Quinton, J. N.; Pachepsky, Y.; van der Putten, W. H.; Bardgett, R. D.; Moolenaar, S.; Mol, G.; Jansen, B.; Fresco, L.O. The significance of soils and soil science towards realization of the United Nations Sustainable Development Goals, *Soil.* **2016**, *2*, 111-128. Doi; 10.5194/soil-2-111-2016.
- (13) Pérez, P.A.; Eugenio, N.P. Status of local soil contamination in Europe: Revision of the indicator "Progress in the management Contaminated Sites in Europe, EUR 29124 EN, Publications Office of the European Union, Luxembourg, **2018**, ISBN 978-92-79-80072-6, Doi:10.2760/093804, JRC107508.
- (14) FAO and ITPS. Global Assessment of the Impact of Plant Protection Products on Soil Functions and Soil Ecosystems FAO, Rome. **2017**, 40 pp
- (15) Covaci, A.; Manirakiza, P.; Schepens, P. Persistent Organochlorine Pollutants in Soils from Belgium, Italy, Greece, and Romania. *Bull. Environ. Contam. Toxicol.* **2002**, *68*, 97-103. Doi: 10.1007/s00128-001-0224-6
- (16) Růžicková, P.; Klánová, J.; Cupr, P.; Lammel, G.; Holoubek, I. An assessment of air-soil exchange of polychlorinated biphenyls and organochlorine pesticides across central and southern Europe. *Environ Sci Technol.* **2008**, *42*(1), 179-85.
- (17) Orton, T.G.; Saby, N.P.A.; Arrouays, D.; Jolivet, C.C.; Villanneau, E.J.; Marchant, B.P. et al. Spatial distribution of Lindane concentration in topsoil across France *Sci. Total Environ.* **2013**, *443*, 338-350.
- (18) Masia, A.; Vasquez, K.; Campo, J.; Pico, Y. Assessment of two extraction methods to determine pesticides in soils, sediments and sludges. Application to the Turia River Basin *J. Chromatogr. A.* **2015**, *1378*, 19-31.
- (19) Pose-Juan, E. Sanchez-Martin, M.J. Andrades, M.S. Rodriguez-Cruz, M.S. Herrero-Hernandez E. Pesticide residues in vineyard soils from Spain: spatial and temporal distributions. *Sci. Total Environ.* **2015**, *514*, 351-358.

- (20)Qu, C.K.; Albanese, S.; Chen, W.; Lima, A.; Doherty, A.L.; Piccolo, A. et al. The status of organochlorine pesticide contamination in the soils of the Campanian Plain, southern Italy, and correlations with soil properties and cancer risk *Environ. Pollut.* **2016**, *216*, 500-511.
- (21)Chiaia-Hernández A.C.; Keller, A.; Waechter, D.; Steinlin, C.; Camenzuli, L.; Hollender J., et al. Long-term persistence of pesticides and tps in archived agricultural soil samples and comparison with pesticide application. *Environ. Sci. Technol.* **2017**, *51*, 10642-10651.
- (22)Ockleford, C.; Adriaanse, P.; Berny, P. Brock, T. Duquesne, S. Grilli, S. et al. Scientific opinion addressing the state of the science on risk assessment of plant protection products for in-soil organisms. *EFSA J.* **2017**, *15*.
- (23)FRN. Federal Republic of Nigeria, Official Gazette, Legal Notice on publication of the details of the Breakdown of the National and State Provisional total, 2006 Census. The Federal Republic of Nigeria Official Gazette, Government Notice No. 21, Lagos. **2007**, *94*, 1-26.
- (24)Martin R.; Samuel F. et al. Interventions to reduce pesticide exposure from the agricultural sector in Africa: a workshop report. *Int. J. Environ. Res. Public Health.* **2022**, *19*, 8973.
- (25)Beyuo, J.; Sackey, L.N.A.; Yeboah, C. et al. The implications of pesticide residue in food crops on human health: a critical review. *Discov Agric.* **2024**, *2*, 123.
- (26)Wondimu, K.T.; Geletu, A.K.; Kedir, W.M. Recent developments in monitoring of organophosphorus pesticides in food samples. *Journal of Agriculture and Food Research.* **2025**, *19*, 101709.
- (27)Liangliang, T.; Dongmei, H.; Yongfu, S.; Feng, H.; Yuan, W.; Hongli, Y.; Huijuan, Y. Method for the Analysis of 7 Indicator Polychlorinated Biphenyls (PCBs) and 13 Organochlorine Pesticide Residues in Sediment by Gas Chromatography (GC). *Earth Environ. Sci.* **2019**, *237*, 022053.
- (28)Indrajit, S.; Samir, V. Environmental Analysis, Analysis of Pesticide residues in drinking water as per bureau of indian standards using the agilent 7000 gc/ms/ms with pesticides analyzer; Solutions for Your Analytical Business Markets and Applications Programs. EU: Agilent Technologies, Inc. **2006**.
- (29)Jona, T.; Yargeau, V. Multiresidue method for the fast and efficient analysis of current-use pesticides in streambed sediments using pressurized liquid extraction. *Science of the Total Environment.* **2024**, *906*, 167703.
- (30)Vincent, K.B.; Nathaniel, O.B.; Lawrence, S.B.; Samuel, A. Human Risk Assessment of Organochlorine Pesticide Residues in Vegetables from Kumasi, Ghana. *J Chem.* **2018**. Doi: 10.1155/2018/3269065.
- (31)Han, Y.; Mo, R.; Yuan, X.; Zhong, D.; Tang, F.; Ye, C; Liu, Y. Pesticide residues in nut-planted soils of China and their relationship between nut/soil. *Chemosphere.* **2017**. Doi:10.1016/j.chemosphere.2017.03.138.
- (32)Chaiyarat, R.; Sookjam, C.; Eiam-Ampai, K. Organochlorine pesticide levels in the food web in rice paddies of Bueng Boraphet wetland, Thailand. *Environ Monit Assess.* **2015**, *187*, 230-240.